



**ISSN: 2454-9940**



**INTERNATIONAL JOURNAL OF APPLIED  
SCIENCE ENGINEERING AND MANAGEMENT**

**E-Mail :**  
**editor.ijasem@gmail.com**  
**editor@ijasem.org**

**[www.ijasem.org](http://www.ijasem.org)**

## Parametric testing statistical hypotheses for fuzzy random variables

Kandikonda Sreekanth<sup>1</sup>, Pothapragada Himabindu<sup>2</sup>, N.Koteswar Rao<sup>3</sup>, Ramarao Tata<sup>4</sup>, Venkateswarrao Deshabona<sup>5</sup>,  
Assistant Professor<sup>1,2,3,4,5</sup>,

Mail Id : [srikanthkandikonda@gmail.com](mailto:srikanthkandikonda@gmail.com), Mail Id : [bindu68287@gmail.com](mailto:bindu68287@gmail.com),

Mail id: [kotichowdaryn@gmail.com](mailto:kotichowdaryn@gmail.com), Mail Id: [ramaraotata12@gmail.com](mailto:ramaraotata12@gmail.com), Mail Id: [venkatdeshaboina69@gmail.com](mailto:venkatdeshaboina69@gmail.com),

Department of H & S,

Swarna Bharati Institute of Science and Technology (SBIT),

Pakabanda Street, Khammam TS, India-507002.

### Abstract:

In this paper, a method is proposed for testing statistical hypotheses about the fuzzy parameter of the underlying parametric population. In this approach, using definition of fuzzy random variables, the concept of the power of test and p value is extended to the fuzzy power and fuzzy p value. To do this, the concepts of fuzzy p value have been defined using the  $\alpha$ -optimistic values of the fuzzy observations and fuzzy parameters. This paper also develops the concepts of fuzzy type-I, fuzzy type-II errors and fuzzy power for the proposed hypothesis tests. To make decision as a fuzzy test, a well-known index is employed to compare the observed fuzzy p value and a given significance value. The result provides a fuzzy test function which leads to some degrees to accept or to reject the null hypothesis. As an application of the proposed method, we focus on the normal fuzzy random variable to investigate hypotheses about the related fuzzy parameters. An applied example is provided throughout the paper clarifying the discussions made in this paper.

1 Introduction The purpose of statistical inference is to draw conclusions about a population on the basis of data obtained from a sample of that population. Hypothesis testing is the process used to evaluate the strength of evidence from the sample and provides a framework for making decisions related to the

population, i.e., it provides a method for understanding how reliably one can extrapolate observed findings in a sample under study to the larger population from which the sample was drawn. The investigator formulates a specific hypothesis, evaluates data from the sample, and uses these data to decide whether they support the specific hypothesis. The classical parametric

approaches usually depend on certain basic assumptions about the underlying population such as: crisp observations, exact parameters, crisp hypotheses, and crisp possible decisions. In practical studies, however, it is frequently difficult to assume that the parameter, for which the distribution of a random variable is determined, has a precise value or the value of the random variable is recorded as a precise value or the hypotheses of interest are presented as exact relations, and so on. Therefore, to achieve suitable testing statistical methods dealing with imprecise information, we need to model the imprecise information and extend the usual approaches to imprecise environments. Since its introduction by Zadeh (1965), fuzzy set theory has been developed and applied in some statistical contexts to deal with uncertainty conditions such as above situations. Specially, the topic of testing statistical hypotheses in fuzzy environments has extensively been studied. Below is a brief review of some studies relevant to the present

work. Arnold (1996, 1998) presented an approach for testing fuzzily formulated hypotheses based on crisp data, in which he proposed and considered generalized definitions of the probabilities of the errors of type-I and type-II. Viertl (2006, 2011) used the extension principle to obtain the generalized estimators for a crisp parameter based on fuzzy data. He also developed some other statistical inferences for the crisp parameter, such as generalized confidence intervals and p value, based on fuzzy data. Taheri and Behboodian (1999) formulated the problem of testing fuzzy hypotheses when the observations are crisp. They presented some definitions for the probabilities of type-I and type-II errors, and proved an extended version of the Neyman–Pearson Lemma. Their approach has been extended by Torabi et al. (2006) to the case in which the data are fuzzy, too. Taheri and Behboodian (2001) also studied the problem of testing hypotheses from a Bayesian point of view when the observations are ordinary and the hypotheses are fuzzy. Taheri and Arefi (2009) presented an approach to the problem of testing fuzzy hypotheses, based on the so-called fuzzy critical regions. Grzegorzewski (2000) suggested some fuzzy tests for crisp hypotheses concerning an unknown parameter of a population using fuzzy random variables (FRVs). Montenegro et al. (2001, 2004), using a generalized metric for fuzzy numbers, proposed a method to test hypotheses about the fuzzy mean of a FRV in one and two populations settings. Gonzalez-Rodríguez et al. (2006) extended a one-sample bootstrap method of testing about the mean of a general fuzzy random variable. Gil et al. (2006) introduced a bootstrap approach to the multiple-sample test of means for imprecisely valued sample data. Chachi and Taheri (2011) introduced a new approach to construct fuzzy

confidence intervals for the fuzzy mean of a FRV. Filzmoser and Viertl (2004) and Parchami et al. (2010) presented p value-based approaches to the problem of testing hypothesis, when the available data or the hypotheses of interest are fuzzy, respectively. Hryniewicz (2006b) investigated the concept of p value in a possibilistic context in which the concept of p value is generalized for the case of imprecisely defined statistical hypotheses and vague statistical data. On the other hand, there have been some studies on non-parametric statistical testing hypotheses in fuzzy environment. Concerning the purposes of this paper, let us briefly review some of the literature on this topic. Kahraman et al. (2004) proposed some algorithms for fuzzy non-parametric rank-sum tests based on fuzzy random variables. Grzegorzewski (1998) introduced a method to estimate the median of a population using fuzzy random variables. He (Grzegorzewski 2004) also demonstrated a straightforward generalization of some classical non-parametric tests for fuzzy random variables based on a metric in the space of fuzzy numbers. He also (Grzegorzewski 2005, 2009) studied some non-parametric median fuzzy tests for fuzzy observations showing a degree of possibility and a degree of necessity (Dubois and Prade 1983) for evaluating the underlying hypotheses. In addition, he (Grzegorzewski 2008) proposed a modification of the classical sign test to cope with fuzzy data which was so-called bi-robust test, i.e., a test which is both distribution free and which does not depend so heavily on the shape of the membership functions used for modeling fuzzy data. Dencœux et al. (2005), using a fuzzy partial ordering on closed intervals, extended the non-parametric ranksum tests based on fuzzy data. For evaluating the hypotheses of interest at a crisp or a fuzzy significance level, they

employed the concepts of fuzzy p value and degree of rejection of the null hypothesis quantified by a degree of possibility and a degree of necessity. Hryniewicz (2006a) investigated the fuzzy version of the Goodman–Kruskal  $\gamma$ -statistic described by ordered categorical data. Lin et al. (2010) considered the problem of two-sample Kolmogorov–Smirnov test for continuous fuzzy intervals based on a crisp test statistic. Taheri and Hesamian (2011) introduced a fuzzy version of the Goodman–Kruskal  $\gamma$ -statistic for two-way contingency tables when the observations were crisp, but the categories were described by fuzzy sets. In this approach, a method was also developed for testing of independence in the two-way contingency tables. Taheri and Hesamian (2012) extended the Wilcoxon signed-rank test to the case where the available observations are imprecise and underlying hypotheses are crisp. Hesamian and Chachi (2013) developed the concepts of fuzzy cumulative distribution function and fuzzy empirical cumulative distribution function and investigated the large sample property of the classical empirical cumulative distribution function for fuzzy empirical cumulative distribution function. They proposed a method for developing two-sample Kolmogorov–Smirnov test for the case when the data are observations of fuzzy random variables, and the hypotheses are imprecise. For more on fuzzy statistics including testing hypotheses for imprecise data, see for example Bertoluzza et al. (2002), Buckley (2006), Kruse and Meyer (1987), Nguyen and Wu (2006), Viertl (2011). This paper develops an approach to test hypotheses for an unknown fuzzy parameter based on fuzzy random variables. To do this, we extend the concept of fuzzy power function and fuzzy p value to investigate the hypotheses of interest. Finally, a decision rule is suggested to

accept or reject the null and alternative hypotheses. We also provide a computational procedure and an example to express the proposed method to test statistical hypotheses for a normal FRV. This paper is organized as follows: Section 2 briefly reviews the classical parametric testing hypotheses and some definitions from fuzzy numbers. In the same section, some results about  $\alpha$ -optimistic values of a fuzzy number are derived to reconstruct a so-called definition of fuzzy random variables introduced in Sect. 3. In Sect. 4, one-sided and two-sided hypotheses about a fuzzy parameter of a FRV are defined. The concept of fuzzy power and fuzzy p value for testing hypotheses about a fuzzy parameter of a continuous parametric population is also introduced. Then, the proposed method is applied for testing hypotheses about the fuzzy parameters of a normal FRV in Sect. 4.1. A numerical example is then provided in Sect. 5 to clarify the discussions made in this paper. Section 6 compares the proposed method to the other similar existing methods. Finally, a brief conclusion is provided in Sect. 7. In addition, the proofs of the main results in this paper are provided in Appendix.

## 2 Preliminaries:

Testing statistical hypotheses: the classical approach



Let  $\mathbf{X} = (X_1, \dots, X_n)$  be a random sample, with observed value  $\mathbf{x} = (x_1, \dots, x_n)$ , from a continuous distribution with density function  $f_\theta$  where  $\theta \in \Theta \subseteq \mathbb{R}^p$ ,  $p \geq 1$ . The decision rule to test the null hypothesis  $H_0 : \theta \in \Theta_0$  versus the alternative  $H_1 : \theta \in \Theta_1$  is typically denoted by

$$\varphi(\mathbf{x}) = \begin{cases} 1 & \text{if } T(\mathbf{x}) \in \mathbf{C}_\delta, \\ 0 & \text{if } T(\mathbf{x}) \notin \mathbf{C}_\delta, \end{cases}$$

where  $\mathbf{C}_\delta \subseteq \mathbb{R}$  is the critical region,  $\delta$  is the significance level and  $T(\mathbf{X})$  is a test statistic. The power function of  $\varphi$  is defined by  $\pi_\varphi(\theta) = E_\theta[\varphi(\mathbf{X})] = \mathbf{P}_\theta(T(\mathbf{X}) \in \mathbf{C}_\delta)$ . If that, if the underlying family of distribution functions has the MLR property (Monotone Likelihood Ratio property) then  $\pi_\varphi(\theta) = E_\theta[\varphi(\mathbf{X})]$  is a non-decreasing function of  $\theta$  (for more see Lehmann and Romano 2005; Shao 2003). Finally, at a given significance level  $\delta$ , the hypothesis is completely rejected if and only if  $p$  value  $< \delta$ , where  $p$  value =  $\inf\{\delta \in [0, 1] : T(\mathbf{x}) \in \mathbf{C}_\delta\}$  (Shao 2003). So, to accept or reject the null hypothesis, we have a test as follows

$$\psi(\mathbf{x}) = \begin{cases} 1 & \text{if } p \text{ value} < \delta, \\ 0 & \text{if } p \text{ value} \geq \delta. \end{cases}$$

### Fuzzy numbers

Let  $\mathbb{R}$  be the set of all real numbers. A fuzzy set of  $\mathbb{R}$  is a mapping  $\mu_A : \mathbb{R} \rightarrow [0, 1]$ , which assigns to each  $x \in \mathbb{R}$  a degree of membership  $0 \leq \mu_A(x) \leq 1$ . For each  $\alpha \in (0, 1]$ , the subset  $\{x \in \mathbb{R} \mid \mu_A(x) \geq \alpha\}$  is called the level set or  $\alpha$ -cut of  $A$  and is denoted by  $A[\alpha]$ . The set  $A[0]$  is also defined equal to the closure of  $\{x \in \mathbb{R} \mid \mu_A(x) > 0\}$ . A fuzzy set  $A$  of  $\mathbb{R}$  is called a fuzzy number if it satisfies the following three conditions

1. For each  $\alpha \in [0, 1]$ , the set  $\tilde{A}[\alpha]$  is a compact interval which will be denoted by  $[\tilde{A}_\alpha^L, \tilde{A}_\alpha^U]$ . Here,  $\tilde{A}_\alpha^L = \inf\{x \in \mathbb{R} \mid \mu_{\tilde{A}}(x) \geq \alpha\}$  and  $\tilde{A}_\alpha^U = \sup\{x \in \mathbb{R} \mid \mu_{\tilde{A}}(x) \geq \alpha\}$ .
2. For  $\alpha, \beta \in [0, 1]$ , with  $\alpha < \beta$ ,  $\tilde{A}[\beta] \subseteq \tilde{A}[\alpha]$ .
3. There is a unique real number  $x^* = x_A^* \in \mathbb{R}$ , such that  $\tilde{A}(x^*) = 1$ . Equivalently, the set  $\tilde{A}[1]$  is a singleton.

The set of all fuzzy numbers is denoted by  $\mathcal{F}(\mathbb{R})$ . Moreover, we denote by  $\text{Fc}(\mathbb{R})$  the set of all fuzzy numbers with continuous membership function.

**Lemma 1** (Lee 2005) Let  $\tilde{A} \in \mathcal{F}(\mathbb{R})$ . Then, both the maps  $\alpha \mapsto \tilde{A}_\alpha^L$  and  $\alpha \mapsto \tilde{A}_\alpha^U$  are left continuous on  $[0, 1]$ .

A LR-fuzzy number  $\tilde{A} = (a, a^l, a^r)_{LR}$  where  $a^l, a^r \geq 0$ , is defined as follows

$$\mu_{\tilde{A}}(x) = \begin{cases} L\left(\frac{a-x}{a^l}\right) & a - a^l \leq x \leq a, \\ R\left(\frac{x-a}{a^r}\right) & a < x \leq a + a^r, \end{cases}$$

where  $L$  and  $R$  are continuous and strictly decreasing functions with  $L(0) = R(0) = 1$  and  $L(1) = R(1) = 0$  (Lee 2005). A special type of LR-fuzzy numbers is the so-called triangular fuzzy numbers with the shape functions  $L(x) = R(x) = \max\{0, 1 - |x|\}$ ,  $x \in \mathbb{R}$ . A well-known ordering of fuzzy numbers, used in the sections below for defining the hypotheses of interest is defined as follows:

**Definition 1** (Wu 2005) Let  $\tilde{A}, \tilde{B} \in \mathcal{F}(\mathbb{R})$ , then

1.  $\tilde{A} \succ (\neq) \tilde{B}$ , if  $\tilde{A}_\alpha^L = (\neq) \tilde{B}_\alpha^L$  and  $\tilde{A}_\alpha^U = (\neq) \tilde{B}_\alpha^U$  for any  $\alpha \in [0, 1]$ .
2.  $\tilde{A} \leq (<) \tilde{B}$ , if  $\tilde{A}_\alpha^L \leq (<) \tilde{B}_\alpha^L$  and  $\tilde{A}_\alpha^U \leq (<) \tilde{B}_\alpha^U$  for any  $\alpha \in [0, 1]$ .
3.  $\tilde{A} \geq (>) \tilde{B}$ , if  $\tilde{A}_\alpha^L \geq (>) \tilde{B}_\alpha^L$  and  $\tilde{A}_\alpha^U \geq (>) \tilde{B}_\alpha^U$  for any  $\alpha \in [0, 1]$ .

**$\alpha$ -Optimistic values** In this subsection, we derive some results about the  $\alpha$ -optimistic values of a fuzzy number. We will use these results to reconstruct a definition of fuzzy random variable that we use in the next sections. For a given fuzzy number  $A$ , the credibility of the event  $\{A \geq r\}$  is defined by Liu (2004) as follows:

$$\text{Cr}\{\tilde{A} \geq r\} := \frac{1}{2} \left( \sup_{y \in [r, +\infty)} \mu_{\tilde{A}}(y) + 1 - \sup_{y \in (-\infty, r)} \mu_{\tilde{A}}(y) \right), \quad (1)$$

It is worth noting that: (1)  $\text{Cr}\{A \geq r\} \in [0, 1]$  and (2)  $\text{Cr}\{A \geq r\} = 1 - \text{Cr}\{A < r\}$ . Here, we recall the definition of the  $\alpha$ -optimistic, but with a small change in the structure of the

original definition. For a fuzzy number A and the real number  $\alpha \in [0, 1]$ , the  $\alpha$  optimistic value of A, denoted by  $A_\alpha$ , is rewritten by  $A_\alpha := \sup\{x \in A [0] \mid Cr\{A \geq x\} \geq \alpha\}$ . (2) Remark 1 It is mentioned that, according to the Liu's definition, the  $\alpha$ -optimistic value of the fuzzy number A is defined as follows  $A_\alpha := \sup\{x \in R \mid Cr\{A \geq x\} \geq \alpha\}$ . (3) Therefore, we observe that  $A_0 = \infty$  and  $A_\alpha \in [A_L \alpha, A_R \alpha)$  for  $\alpha \in (0, 1]$ . While, by Eq. (2), each value of  $A_\alpha$  belongs to  $A [0]$  (which is a compact interval, due to the definition of a fuzzy number), for all  $\alpha \in [0, 1]$ . Moreover, it is clear that  $A_\alpha$  is a non-increasing function of  $\alpha \in [0, 1]$  (see Liu 2004, for more details).

#### Numerical example

In this section, we provide a numerical example to clarify the discussions in this paper and give the possible application of the proposed method for normal FRVs. Example 2 (Elsherif et al. 2009) Based on a random sample of the received signal from a target, assume we have fuzzy observations  $x = (x_1, x_2, \dots, X_{25})$  (as shown in Table 1) measured in nanowatt which is approximately normally distributed  $N(\theta, \sigma^2)$ . We wish to test the following hypothesis:

$$\begin{cases} \tilde{H}_0 : \tilde{\theta} \asymp \tilde{\theta}_0 \asymp (1.35, 1.50, 1.80)_T, \\ \tilde{H}_1 : \tilde{\theta} < \tilde{\theta}_0. \end{cases}$$

Applying the procedure proposed in Sect. 4.1, using Remark 6, the fuzzy p value is derived point by point and its membership function is shown in Fig. 1. This membership function can be interpreted as “about 0.075”. Therefore, at level of  $\delta = 0.05$ , the fuzzy test for testing  $H_0 : \theta \sim (1.35, 1.50, 1.80)_T$  is obtained as  $\psi(X) = \left\{ \begin{matrix} 0.905 \\ 0 \\ 0.095 \\ 1 \end{matrix} \right\}$ . So, with respect to the observed fuzzy observations and at significance level of  $\delta = 0.05$ , the null hypothesis  $H_0 : \theta \sim (1.35,$

$1.50, 1.80)_T$  is accepted with a degree of 0.905 and it is rejected with a degree of 0.095. Such a result may be interpreted as “we absolutely tend to accept  $H_0$ ”. In addition, based on Remark 7, the membership of the fuzzy power at  $\theta^* (0.50, 0.70, 0.80)_T$  is shown in Fig. 2 which is interpreted as “about 0.999”.

Now, suppose that we want to test the following hypotheses about  $\sigma^2$ :

$$\begin{cases} \tilde{H}_0 : \tilde{\sigma}^2 \asymp \tilde{\sigma}_0^2 \asymp (0.30, 0.50, 0.70)_T, \\ \tilde{H}_1 : \tilde{\sigma}^2 > \tilde{\sigma}_0^2. \end{cases}$$

Table 1 The observed fuzzy random sample in Example 2

$\tilde{x}_1 = (0.02, 0.10, 0.23)_T$	$\tilde{x}_2 = (0.08, 0.20, 0.37)_T$	$\tilde{x}_3 = (0.15, 0.30, 0.46)_T$
$\tilde{x}_4 = (0.28, 0.40, 0.56)_T$	$\tilde{x}_5 = (0.37, 0.50, 0.65)_T$	$\tilde{x}_6 = (0.42, 0.60, 0.74)_T$
$\tilde{x}_7 = (0.54, 0.70, 0.83)_T$	$\tilde{x}_8 = (0.66, 0.80, 0.93)_T$	$\tilde{x}_9 = (0.72, 0.90, 1.26)_T$
$\tilde{x}_{10} = (0.85, 1.00, 1.15)_T$	$\tilde{x}_{11} = (0.98, 1.10, 1.22)_T$	$\tilde{x}_{12} = (1.05, 1.20, 1.32)_T$
$\tilde{x}_{13} = (1.17, 1.30, 1.48)_T$	$\tilde{x}_{14} = (1.25, 1.40, 1.54)_T$	$\tilde{x}_{15} = (1.32, 1.50, 1.67)_T$
$\tilde{x}_{16} = (1.37, 1.60, 1.78)_T$	$\tilde{x}_{17} = (1.53, 1.70, 1.88)_T$	$\tilde{x}_{18} = (1.62, 1.80, 1.97)_T$
$\tilde{x}_{19} = (1.75, 1.90, 2.08)_T$	$\tilde{x}_{20} = (1.76, 2.00, 2.23)_T$	$\tilde{x}_{21} = (1.93, 2.10, 2.27)_T$
$\tilde{x}_{22} = (2.06, 2.20, 2.35)_T$	$\tilde{x}_{23} = (2.05, 2.30, 2.46)_T$	$\tilde{x}_{24} = (2.21, 2.40, 2.55)_T$
$\tilde{x}_{25} = (2.32, 2.50, 2.60)_T$	-	-




Fig. 1 The membership function of the fuzzy p value and the significance level in Example 2

for testing  $H_0 : \theta \sim (1.35, 1.50, 1.80)T$  against  
 $H_1 : \theta < (1.35, 1.50, 1.80)T$




Fig. 2 The membership function of the fuzzy power at  $\hat{\theta}^* \sim (0.50, 0.70, 0.80)T$  in Example 2




Fig. 3 The membership function of the fuzzy p value and the significance level in Example 2 for testing  $H_0 : \sigma^2 (0.30, 0.50, 0.70)T$  against  $H_1 : \sigma^2 (0.30, 0.50, 0.70)T$

### A comparison study

It should be mentioned that Akbari and Rezaei (2009, 2010) extended the classical approach for testing statistical hypotheses about the (crisp) mean or variance of a normal distribution by introducing some fuzzy numbers to interpret the sentences “larger

than”, “smaller than” or “not equal” for population’s parameters based on imprecise observations. Parchami et al. (2010) defined an extension of fuzzy p value for a crisp random sample of a normal distribution to test imprecisely hypothesis test about the mean. Arefi and Taheri (2011) extended the classical critical region for testing fuzzy hypothesis about the parameters of the classical normal distribution based on fuzzy data. Wu (2005) considered the problem of hypotheses test with fuzzy data. Based on a concept of fuzzy random variable, first he defined the fuzzy mean of the normal distribution. Then, based on a proposed ranking method, he defined the hypotheses about the population’s fuzzy mean. He considered two cases: (1) the variance of the population is known, (2) the variance of the population is unknown. Finally, he introduced a fuzzy test based on the classical critical region. However, for both cases, he assigned a crisp number into the structure of testing hypothesis’s method: for the case (1) the variance of the population is considered as a crisp number and for the case (2) the variance of the fuzzy sample is calculated based on core of the fuzzy data as a crisp number. As we observe, in all above methods, the variance of the normal population is considered as a crisp number. However, for testing fuzzy hypothesis about mean of a normal population, it is reasonable that the variance of the population is also considered as a imprecise value. So, it may be an advantage of our method with respect to abovementioned methods. Geyer and Meeden considered the problem of the classical optimal hypothesis tests for the binomial distribution (in general for discrete data with crisp parameter) using an interval estimation of a binomial proportion (Geyer and Meeden 2005). They introduced the notion of fuzzy confidence intervals by inverting families of

randomized tests. Then, they introduce a notion of fuzzy p value in which it is only a function of the parameters of the model. Finally, they provided a unified description of fuzzy confidence interval, fuzzy p values and fuzzy decision. However, their proposed method is not a fuzzy method in general, since this approach does not consider the imprecise information about the model such as imprecise observations or imprecise hypotheses and it does not lead to a fuzzy decision. Viertl (2011) and Filzmoser and Viertl(2004) also extend a concept of fuzzy p value when the observations are fuzzy and hypotheses are crisp. Arnold (1996, 1998) proposed a method for testing fuzzy hypotheses about the population parameter with crisp data. He provided some definitions for the probability of type-I and type-II errors and presented the best test for the oneparameter exponential family.

As it is observed, in all above-proposed methods for testing statistical hypothesis, at least one of the essential population's information such as data, hypothesis or population's parameters play a crisp role in the structure of testing procedure. While, by restructuring the concept of fuzzy random variable, we involved the population's information including: fuzzy data and fuzzy parameters into the hypothesis test, type-I error (or type-II error) and power of test. Finally, we obtained a fuzzy test to make decision for accepting or rejecting the fuzzy hypothesis of interest about the fuzzy parameters.

## Conclusions

This paper proposes a method for testing hypotheses about the fuzzy parameters of a fuzzy random variable. In this approach, we reconstruct a well-known concept of a fuzzy

random variable using the concept of  $\alpha$ -optimistic values. Then, we extended the concepts of the fuzzy power test and fuzzy p value. Finally, based on the credibility index, to compare the observed fuzzy p value and the crisp significance level, the fuzzy hypothesis of interest can be accepted or rejected with degrees of conviction between 0 and 1. Although we focused on testing hypotheses about the parameters of a normal fuzzy random variable, the proposed method is general and it can be applied for other kinds of fuzzy random variables as well. Moreover, it can be applied for other kind of testing hypothesis such as comparing two independent fuzzy random variables of two populations, the one-way or two-way analysis of variance and nonparametric approaches including testing hypothesis. However, the proposed method can be applied only for fuzzynumbers involved in a problem of statistical hypothesis testing. Moreover, the topic of testing hypothesis for fuzzy parameters of a fuzzy random variable can be extended to the case where the level of significance is given by a fuzzy number, too. To do this, one can easily apply the credibility index for comparing two fuzzy numbers into the decision making.

## Appendix: Proof of the main results

This appendix provides the mathematical proofs of the theoretical results in our paper.



*Proof of Theorem 1* (i) Let  $g_{\tilde{A}} : \mathbb{R} \rightarrow [0, 1]$  be given by

$$g_{\tilde{A}}(x) = \text{Cr}(\tilde{A} \geq x). \text{ Then}$$

$$\tilde{A}_\alpha = \sup\{x \in \tilde{A}[0] \mid g_{\tilde{A}}(x) \geq \alpha\}.$$

First, let  $\alpha \in [0, 0.5)$ . Since

$$\begin{aligned} g_{\tilde{A}}(\tilde{A}_{2\alpha}^U) &= \frac{1}{2} \left( \sup_{y \in [\tilde{A}_{2\alpha}^U, +\infty)} \mu_{\tilde{A}}(y) + 1 - \sup_{y \in (-\infty, \tilde{A}_{2\alpha}^U)} \mu_{\tilde{A}}(y) \right) \\ &= \frac{1}{2}(2\alpha) = \alpha, \end{aligned}$$

we have  $\tilde{A}_\alpha \geq \tilde{A}_{2\alpha}^U$ . On the other hand, for each  $z > \tilde{A}_{2\alpha}^U$ ,  $\sup_{y \in [z, +\infty)} \mu_{\tilde{A}}(y) = \tilde{A}(z)$  and  $\sup_{y \in (-\infty, z)} \mu_{\tilde{A}}(y) = 1$ . Therefore,

$$\begin{aligned} g_{\tilde{A}}(z) &= \text{Cr}(\tilde{A} \geq z) \\ &= \frac{1}{2} \left( \sup_{y \in [z, +\infty)} \mu_{\tilde{A}}(y) + 1 - \sup_{y \in (-\infty, z)} \mu_{\tilde{A}}(y) \right) \\ &= \frac{1}{2} \tilde{A}(z) < \alpha. \end{aligned}$$

Hence,  $\{x \in \mathbb{R} \mid g_{\tilde{A}}(x) \geq \alpha\} \cap (\tilde{A}_{2\alpha}^U, +\infty) = \emptyset$ , which by the previous part implies that  $\tilde{A}_\alpha = \tilde{A}_{2\alpha}^U$ .

Now suppose  $\alpha \in [0.5, 1]$  and let  $x_0 := \sup\{x \leq x^* \mid \mu_{\tilde{A}}(x) \leq 2(1 - \alpha)\}$ . Then,  $\mu_{\tilde{A}}(x) \leq 2(1 - \alpha)$ , for all  $x < x_0$ . Hence,

$$\sup_{y \in [x_0, +\infty)} \mu_{\tilde{A}}(y) = 1 \text{ and } \sup_{y \in (-\infty, x_0)} \mu_{\tilde{A}}(y) \leq 2(1 - \alpha).$$

Therefore,  $g_{\tilde{A}}(x_0) \geq \alpha$  which implies that  $\phi_{\tilde{A}}(\alpha) \geq x_0$ . On the other hand, for  $x > x_0$ , since  $\mu_{\tilde{A}}(x) > 2(1 - \alpha)$ ,

we have

$$\begin{aligned} g_{\tilde{A}}(x) &= \frac{1}{2} \left( \sup_{y \in [x, +\infty)} \mu_{\tilde{A}}(y) + 1 - \sup_{y \in (-\infty, x)} \mu_{\tilde{A}}(y) \right) \\ &\geq \frac{1}{2}(1 + 1 - \mu_{\tilde{A}}(x)) < \alpha. \end{aligned}$$

Hence,  $\phi_{\tilde{A}}(\alpha) = x_0$ . Note that, according to what has been proved,  $\phi_{\tilde{A}}(0.5) = \sup\{x \leq x^* \mid \mu_{\tilde{A}}(x) \leq 1\} = x^* = \tilde{A}_{2(0.5)}^U$ . Thus, the law of  $\phi_{\tilde{A}}$  can be written in the following form.

$$\forall \alpha \in [0, 1], \quad \phi_{\tilde{A}}(\alpha) = \begin{cases} \tilde{A}_{2\alpha}^U & 0 \leq \alpha \leq 0.5, \\ \sup\{x \leq x^* \mid \mu_{\tilde{A}}(x) \leq 2(1 - \alpha)\} & 0.5 \leq \alpha \leq 1. \end{cases}$$

(ii) It is clear that  $\phi_A$  is decreasing on  $[0, 1]$ . By the previous part, since for each  $\alpha \in [0, 0.5]$ ,  $\phi_A(\alpha) = A \cup 2\alpha$ , using Lemma 1,  $\phi_A$  is left continuous on  $[0, 0.5]$ . For  $\alpha \in (0.5, 1]$ , suppose  $\{\alpha_n\}_{n \in \mathbb{N}}$  is an increasing sequence in  $(0.5, 1]$  with  $\alpha_n \rightarrow \alpha$ . First, suppose that  $\phi_A(\alpha) = x^*$ . Then, for each  $x < x^*$ ,  $\mu_A(x) \leq 2(1 - \alpha) \leq 2(1 - \alpha_n)$ . Hence  $\phi_A(\alpha_n) \geq x^*$ , for all  $n \in \mathbb{N}$ . On the other hand, since  $\alpha_n > 0.5$ , we have  $\phi_A(\alpha_n) \leq x^*$ . Combining the two arguments, we obtain  $\phi_A(\alpha_n) = x^*$ , for all  $n \in \mathbb{N}$ . Hence, in this case  $\phi_A(\alpha_n) = x^* \rightarrow \phi_A(\alpha) = x^*$ . Now suppose  $\phi_A(\alpha) < x^*$ . Then, for each  $y \in \mathbb{R}$  with  $\phi_A(\alpha) < y < x^*$ ,  $\mu_A(y) > 2(1 - \alpha)$ . Therefore,  $\mu_A(y) > 2(1 - \alpha_n)$ , for large values of  $n \in \mathbb{N}$ . This implies that  $\phi_A(\alpha_n) \leq y$ . Hence  $\phi_A(\alpha_n) \rightarrow \phi_A(\alpha)$ . This completes the proof of the left continuity of  $\phi_A$  on  $(0.5, 1]$ .

*Proof of Lemma 2* First suppose  $\alpha_0 \in (0.5, 1]$  is a point of continuity of  $\psi_A$ . Let  $x_0 := \phi_A(\alpha_0)$ . Then  $A \cup L 2(1 - \alpha_0) \leq x_0$ . If  $A \cup L 2(1 - \alpha_0) < x_0$  then  $\mu_A(x) = 2(1 - \alpha_0)$ , for all  $x \in [A \cup L 2(1 - \alpha_0), x_0)$ . In this case for each  $\alpha > 2(1 - \alpha_0)$ , we have  $A \cup L 2(1 - \alpha) \geq x_0$  which contradicts the assumption of continuity of  $\psi_A$  at  $\alpha_0$ . Hence

$\psi_A(\alpha_0) = A L 2(1-\alpha_0) = x_0 = \varphi_A(\alpha_0)$ .

Suppose, on the contrary, that  $\alpha_0$  is a point of discontinuity of  $\psi_A$ . Then, there  $0 > 0$  such that for each  $n \in \mathbb{N}$ , one can find  $\alpha_n \in (\alpha_0 - 1/n, \alpha_0)$  with  $\psi_A(\alpha_n) > \psi_A(\alpha_0) + 0$ . Without loss of generality, we may assume that  $\{\alpha_n\}_{n \in \mathbb{N}}$  is an increasing sequence. Since  $\psi_A$  is a monotonic function, there exists an increasing sequence  $\{\alpha_n\}_{n \in \mathbb{N}}$  converging to  $\alpha_0$  such that  $\alpha_n \leq \alpha_0$ , for each  $n \in \mathbb{N}$ , and  $\psi_A$  is continuous on the set  $\{\alpha_n \mid n \in \mathbb{N}\}$ . Let also  $\{\beta_n\}_{n \in \mathbb{N}}$  be a decreasing sequence converging to  $\alpha_0$  which consist of continuity points of  $\psi_A$ .

Then  $\varphi_A(\alpha_n) = \psi_A(\alpha_n) \geq \psi_A(\alpha_n) \geq \psi_A(\alpha_0) + 0 > \psi_A(\alpha_0) \geq \psi_A(\beta_n) = \varphi_A(\beta_n)$ ,

for each  $n \in \mathbb{N}$ . Hence

$\forall n \in \mathbb{N}, \varphi_A(\alpha_n) - \varphi_A(\beta_n) > 0$ ,

which implies that  $\varphi_A$  is also discontinuous at this point.

## References :

Akbari MG, Rezaei A (2009) Statistical inference about the variance of fuzzy random variables. *Sankhy - Indian J Stat* 71.B(Part 2):206–221  
 Akbari MG, Rezaei A (2010) Bootstrap testing fuzzy hypotheses and observations on fuzzy statistic. *Expert SystAppl* 37:5782–5787  
 Arefi M, Taheri SM (2011) Testing fuzzy hypotheses using fuzzy data based on fuzzy test statistic. *J Uncertain Syst* 5:45–61  
 Arnold BF (1996) An approach to fuzzy hypothesis testing. *Metrika* 44(1):119–126  
 Arnold BF (1998) Testing fuzzy hypothesis with crisp data. *Fuzzy Sets Syst* 94(3):323–333  
 Bertoluzza C, GilMA, Ralescu DA (2002) Statistical modeling, analysis and management of fuzzy data. *Physica-Verlag, Heidelberg*  
 Buckley JJ (2006) *Fuzzy statistics; studies in fuzziness and soft*

*computing*. Springer, Berlin  
 Chachi J, Taheri SM (2011) Fuzzy confidence intervals for mean of Gaussian fuzzy random variables. *Expert SystAppl* 38:5240–5244  
 Dencœux T, Masson MH, Herbert PH (2005) Non-parametric rank-based statistics and significance tests for fuzzy data. *Fuzzy Sets Syst* 153:1–28  
 Dubois D, Prade H (1983) Ranking of fuzzy numbers in the setting of possibility theory. *InfSci* 30:183–224  
 Elsharif AK, Abbady FM, Abdelhamid GM (2009) An application of fuzzy hypotheses testing in radar detection. In: 13th international conference on aerospace science and aviation technology, ASAT13, pp 1–9  
 Filzmoser P, Viertl R (2004) Testing hypotheses with fuzzy data: the fuzzy p-value. *Metrika* 59:21–29  
 Gil MA, Montenegro M, Rodríguez G, Colubi A, Casals MR (2006) Bootstrap approach to the multi-sample test of means with imprecise data'. *Comput Stat Data Anal* 51:148–162  
 Gonzalez-Rodríguez G, Montenegro M, Colubi A, GilMA (2006) Bootstrap techniques and fuzzy random variables: Synergy in hypothesis testing with fuzzy data. *Fuzzy Sets Syst* 157:2608–2613  
 Grzegorzewski P (1998) Statistical inference about the median from vague data. *Control Cybern* 27:447–464  
 Grzegorzewski P (2000) Testing statistical hypotheses with vague data. *Fuzzy Sets Syst* 11:501–510  
 Grzegorzewski P (2004) Distribution-free tests for vague data. In: Lopez-Diaz M et al (eds) *Soft methodology and random information systems*. Springer, Heidelberg, pp 495–502  
 Grzegorzewski P (2005) Two-sample median test for vague data. In: Proceedings of the 4th conference European society for fuzzy logic and technology-Eusflat, Barcelona, pp 621–626  
 Grzegorzewski P (2008) A bi-robust test for vague data. In: Magdalena L, Ojeda-Aciego M, Verdegay JL (eds) *Proceedings of the 12th international conference on information*

- processing and management of uncertainty in knowledge-based systems, IPMU2008, Spain, Torremolinos (Malaga), June 22–27, pp 138–144 Grzegorzewski P (2009) K-sample median test for vague data. *Int J IntellSyst* 24:529–539
- Geyer C, Meeden G (2005) Fuzzy and randomized confidence intervals and p-values. *Stat Sci* 20:358–366 Hesamian G, Taheri SM (2013) Fuzzy empirical distribution: properties and applications. *Kybernetika* 49:962–982
- Hesamian G, Chachi J (2013) Two-sample KolmogorovSmirnov fuzzy test for fuzzy random variables. *Stat Pap*.doi:10.1007/s00362-013-0566-2 Hryniewicz O (2006a) Goodman-Kruskal measure of dependence for fuzzy ordered categorical data. *Comput Stat Data Anal* 51:323–334 Hryniewicz O (2006b) Possibilistic decisions and fuzzy statistical tests. *Fuzzy Sets Syst* 157:2665–2673
- Kahraman C, Bozdogan CF, Ruan D (2004) Fuzzy sets approaches to statistical parametric and non-parametric tests. *Int J IntellSyst* 19:1069–1078 Kruse R, Meyer KD (1987) *Statistics with vague data*. D. Riedel Publishing, New York Kwakernaak H (1978) Fuzzy random variables, part I: definitions and theorems. *InfSci* 19:1–15 Kwakernaak H (1979) Fuzzy random variables, part II: algorithms and examples for the discrete case. *InfSci* 17:253–278 Lee KH (2005) *First course on fuzzy theory and applications*. Springer, Berlin Lehmann EL, Romano JP (2005) *Testing statistical hypotheses*, 3rd edn. Springer, New York Liu B (2004) *Uncertainty theory*. Springer, Berlin Lin P, Wu B, Watada J (2010) Kolmogorov–Smirnov two sample test with continuous fuzzy data. *AdvIntell Soft Comput* 68:175–186
- Montenegro M, Casals MR, Lubiano MA, Gil MA (2001) Two-sample hypothesis tests of means of a fuzzy random variable. *InfSci* 133:89–100 Montenegro M, Colubi A, Casals MR, Gil MA (2004) Asymptotic and Bootstrap techniques for testing the expected value of a fuzzy random variable. *Metrika* 59:31–49
- Nguyen H, Wu B (2006) *Fundamentals of statistics with fuzzy data*. Springer, Netherlands Parchami A, Taheri SM, Mashinchi M (2010) Fuzzy p-value in testing fuzzy hypotheses with crisp data. *Stat Pap* 51:209–226 Shao J (2003) *Mathematical statistics*. Springer, New York Taheri SM, Arefi M (2009) Testing fuzzy hypotheses based on fuzzy test statistic. *Soft Comput* 13:617–625 Taheri SM, Behboodian J (1999) Neyman–Pearson lemma for fuzzy hypothesis testing. *Metrika* 49:3–17 Taheri SM, Behboodian J (2001) A Bayesian approach to fuzzy hypotheses testing. *Fuzzy Sets Syst* 123:39–48 Taheri SM, Hesamian G (2011) Goodman–Kruskal measure of association for fuzzy-categorized variables. *Kybernetika* 47:110–122 Taheri SM, Hesamian G (2012) A generalization of the Wilcoxon signed-rank test and its applications. *Stat Pap* 54:457–470
- Torabi H, Behboodian J, Taheri SM (2006) Neyman–Pearson lemma for fuzzy hypothesis testing with vague data. *Metrika* 64:289–304 Viertl R (2006) Univariate statistical analysis with fuzzy data. *Comput Stat Data Anal* 51:133–147 Viertl R (2011) *Statistical methods for fuzzy data*. Wiley, Chichester Wu HC (2005) *Statistical hypotheses testing for fuzzy data*. *InfSci* 175:30–57 Zadeh LA (1965) Fuzzy sets. *Inf Control* 8:338–353.

